arXiv:math/0301139v2 [math.DG] 4 May 2003

Relative isoperimetric inequalities for Lagrangian disks

Sung Ho Wang

Department of Mathematics

Postech

Pohang, Korea 790 - 784

email: **wang@postech.ac.kr**

Given a closed curve $C \subset \mathbb{C}^n$ with zero symplectic area, Allcock showed that C bounds an isotropic disk l with

$$Area(l) \,\leq\, \mu\, Length^2(C)$$

for some universal constant μ by constructing an explicit isotropic deformation of C to a single point. In this note, we wish to prove a relative analogue of this result. Our interest comes from the application of the relative isoperimetric inequality to the regularity of an area minimizing Lagrangian surface [Wa].

Let $\varpi = dx^1 \wedge dy^1 + dx^2 \wedge dy^2$ be the standard symplectic form on $\mathbb{C}^2$ with coordinates $z^k = x^k + i\, y^k$, $k = 1,\, 2$, and let

$$\eta = \frac{1}{2} \sum_k (x^k\, dy^k - \, y^k dx^k).$$

Suppose $\Gamma \subset \mathbb{C}^2$ is a Lagrangian plane through the origin, or a union of two Lagrangian planes that intersect transversally at the origin. For a connected curve segment

$C \subset \mathbb{C}^2$ with boundary points in Γ, consider the integral

$$I(C) = \int_C \eta.$$

Since η vanishes on Γ, $I(C)$ is invariant under isotropic deformation of C with end points in Γ. Moreover, such a curve C bounds a Lagrangian half disk or a wedge if and only if $I(C) = 0$.

On the other hand, suppose Γ is a complex plane through the origin. For any curve C with boundary points in Γ, let $\hat{C} = C \cup r_\Gamma(C)$ be the loop where r_Γ is the reflection across Γ. Then $I(\hat{C}) = 0$, and $\hat{C}$ always bounds a Lagrangian disk.

The following relative isoperimetric inequality can be considered as a quantitative analysis of these observations. Let D be the closed unit disk in the plane, and $D^- = \{\, (x, y) \in D \,|\, x \leq 0 \,\}$. Denote $S^- = \{\, (x, y) \in \partial D^- \,|\, x^2 + y^2 = 1\}$, and $I_y = \{\, (x, y) \in \partial D^- \,|\, x = 0\}$.

Theorem 1 *Let $\Gamma \subset \mathbb{C}^2$ be either a Lagrangian plane, or a complex plane, or a union of any two of Lagrangian or complex planes that intersect transversally at the origin. There exists a constant $\mu(\Gamma)$ such that for any $W^{1,2}$ curve $C : S^- \to \mathbb{C}^2$ which bounds a $W^{1,2}$ Lagrangian half disk(or a wedge respectively in case Γ consists of two planes) $l_0 : D^- \to \mathbb{C}^2$ with $l_0(I_y) \subset \Gamma$ and $l_0|_{S^-} = C$, there exists a $W^{1,2}$ Lagrangian half disk(wedge) $l : D^- \to \mathbb{C}^2$ spanning Γ and C such that*

$$Area(l(D^-)) \leq \mu \, Length^2(C).$$

If Γ contains a complex plane, any curve C with boundary in Γ always bounds a Lagrangian half disk or a wedge with boundary in Γ.

Proof.

1. Case Γ is a Lagrangian plane, or a union of two transversal Lagrangian planes: Suppose Γ is a Lagrangian plane. Let $\gamma \subset \Gamma$ be the straight line between the end points

of C. Since Γ is Lagrangian and C bounds a Lagrangian disk, the symplectic area of the loop $C \cup \gamma$ is zero, and it bounds a Lagrangian disk. The proposition follows from the Gromov-Allcock isoperimetric inequality for $C \cup \gamma$ as $Length(\gamma) \leq Length(C)$. The case of two transversal Lagrangian planes follows similarly.

2. Case Γ is a complex plane : Let z^1, z^2 be the standard coordinates of $\mathbb{C}^2$, and assume Γ is a z^1- plane. Let

$$\alpha(s) = \alpha^1(s) + \alpha^2(s), \ \ 0 \leq \ s \ \leq 1$$

be a parametrization of C with $\alpha(1) = 0$, and each α^i curve lies in z^i-plane. Note α^2 is a closed curve. By Lemma 2.1 [Al], there exists an isotropic homotopy of α to a curve β which first transverses α^1, and then α^2.

Lemma 2.2 [Al] gives an isotropic homotopy of α^2 to a closed curve in z^1-plane, and β is isotropic homotopic to a curve β' in z^1-plane.

The total area of deformation $\alpha \to \beta$ is bounded by $Length^2(C)$, and area of $\beta \to \beta'$ bounded by $\pi \, Length^2(C)$.

3. Case Γ is a union of two transversal complex planes, or a union of a complex and a Lagrangian planes: Let p_1, p_2 be the end points of C, and $L_i = |p_i|$. Then one easily verifies $L_i \leq Length(C) \csc\theta$, where $\theta > 0$ is the minimum angle between the two planes. The theorem follows from Case 2 by taking a straight line from the origin to an end point of C. $\square$

Let X be a Kähler surface. A finite distinct ordered set $\{P_1, P_2, ... P_m = P_0\} = \Gamma \subset X$ is called a *geodesic polygon* if Γ is connected and each *face* P_i is either a complex curve which is a fixed point locus of a Kähler involution r_i, or a Lagrangian surface which is a fixed point locus of an anti-Kähler involution r_i of X. Γ is called *rational* if the group generated by $\{r_i, r_{i+1}\}$ is finite for each $i = 0, 1, ... m-1$.

Let D be the unit disk in the plane, and $W^{1,2}(D, X)$ be the Sobolev space of maps with square integrable first derivative. An element $l \in W^{1,2}(D, X)$ with $\partial l : \partial D \to \Gamma$ is called *monotone* if there exists a partition of ∂D on which ∂l is a monotone map into Γ, [Wa] for detail.

Corollary 1 *Let* $\Gamma \subset X$ *be a rational geodesic polygon in a Kähler surafce. Suppose* $l \in W^{1,2}(D, X)$ *with* $\partial l(\partial D) \subset \Gamma$ *is a weakly conformal, monotone, weakly Lagrangian map that is minimizing in its relative homotopy class. Then* l *is Hölder continuous on* D *up to boundary.*

Proof. The interior regularity is proved in Theorem 2.8 in [ScW] by applying Morrey's argument. Suppose $q \in \partial D$ is a *vertex*, and consider a small extension $\hat{l}$ of l in a neighborhood of q by finite successive reflections across the corresponding faces of Γ, which is defined since l is monotone and Γ is rational. The corollary follows from the relative isoperimetric inequality and the interior regularity applied to $\hat{l}$. Regularity along the edges follows similarly. □

References

[Al] D. Allcock, *An Isoperimetric Inequality for the Heisenberg Groups*, Geom. Funct. Anal. Vol 8, No 2 (1998), 219–233.

[Wa] S. H. Wang, *Compact special Legendrian surfaces in* S^5, arXiv. math. DG/0211439

Sung Ho Wang
Department of Mathematics
Postech
Pohang, Korea 790-784
email: **wang@postech.ac.kr**